

MEDICATIONS FOR OPIOID USE DISORDER (MOUD)

MOUD is an effective, clinically proven treatment that helps people in their recovery. People recover differently - one way people recover is with MOUD.

MOUD is the use of FDA approved medications, typically alongside counseling and behavioral therapies, to provide a "whole-patient" approach to the treatment of opioid use disorders (OUD).

Research shows that for many, a combination of medication and therapy can successfully treat OUD, sustain recovery, and prevent or reduce opioid overdose.

TREATMENT

PROCESS

EFFECTS

DISPENSING

FDA APPROVAL

METHADONE

- Medication binds to opioid receptors
- Relieves cravings and withdrawal symptoms
- On site daily at Opioid Treatment Programs
- 1972

BUPRENORPHINE

- Medication binds to opioid receptors but does not completely activate them
- Reduces withdrawal symptoms and cravings
- Prescribing healthcare provider
- 2002

NALTREXONE

- Medication blocks activation of opioid receptors
- Prevents all opioids from causing euphoria
- Prescribing healthcare provider
- 2006



MOUD treatment is associated with a 76% reduction in overdose at 3 months.



MOUD treatment is associated with a 59% reduction in overdose at 12 months.

LEARN MORE ABOUT UNSHAME NC HERE!

